

Economics 204

Lecture 9—Thursday, August 6, 2009

Revised 8/6/09, revisions indicated by ** and Sticky

Notes

Section 3.3 Supplement, Quotient Vector Spaces (not in de la Fuente):

Definition 1 Given a vector space X and a vector subspace W of X , define an equivalence relation by

$$x \sim y \Leftrightarrow x - y \in W$$

Form a new vector space X/W : the set of vectors is

$$\{[x] : x \in X\}$$

where $[x]$ denotes the equivalence class of x with respect to \sim . Note that the vectors are *sets*; this is a little weird at first, but Define

$$\begin{aligned}[x] + [y] &= [x + y] \\ \alpha[x] &= [\alpha x]\end{aligned}$$

You should check on your own that \sim is an equivalence relation and that vector addition and scalar multiplication are well-defined, i.e.

$$\begin{aligned}[x] = [x'], [y] = [y'] &\Rightarrow [x + y] = [x' + y'] \\ [x] = [x'], \alpha \in F &\Rightarrow [\alpha x] = [\alpha x']\end{aligned}$$

Theorem 2 *If $\dim X < \infty$, then*

$$\dim(X/W) = \dim X - \dim W$$

Theorem 3 *Let $T \in L(X, Y)$. Then $\text{Im } T$ is isomorphic to $X/\ker T$.*

Proof: If $\dim X < \infty$, then $\dim X/\ker T = \dim X - \dim \ker T$ (from the previous theorem) = $\text{Rank } T$ (from Theorem 11 in yesterday's lecture) = $\dim \text{Im } T$, so $X/\ker T$ is isomorphic to $\text{Im } T$. We shall prove that it is true in general, and that the isomorphism is natural. Define

$$\tilde{T}([x]) = T(x)$$

We need to check that this is well-defined.

$$\begin{aligned} [x] = [x'] &\Rightarrow x \sim x' \\ &\Rightarrow x - x' \in \ker T \\ &\Rightarrow T(x - x') = 0 \\ &\Rightarrow T(x) = T(x') \end{aligned}$$

so \tilde{T} is well-defined. Clearly, $\tilde{T} : X/\ker T \rightarrow \text{Im } T$. It is easy to check that \tilde{T} is linear, so $\tilde{T} \in L(X/\ker T, \text{Im } T)$.

$$\begin{aligned} \tilde{T}([x]) = \tilde{T}([y]) &\Rightarrow T(x) = T(y) \\ &\Rightarrow T(x - y) = 0 \\ &\Rightarrow x - y \in \ker T \\ &\Rightarrow x \sim y \\ &\Rightarrow [x] = [y] \end{aligned}$$

so \tilde{T} is one-to-one.

$$\begin{aligned} y \in \text{Im } T &\Rightarrow \exists_{x \in X} T(x) = y \\ &\Rightarrow \tilde{T}([x]) = y \end{aligned}$$

so \tilde{T} is onto, hence \tilde{T} is an isomorphism. ■

Example: Consider $T \in L(\mathbf{R}^3, \mathbf{R}^2)$ defined by

$$T(x, y, z) = (x, y)$$

Then $\ker T = \{(x, y, z) \in \mathbf{R}^3 : x = y = 0\}$ is the z -axis. Given (x, y, z) , the equivalence class $[(x, y, z)]$ is just the line through $(x, y, 0)$ parallel to the z -axis. $\tilde{T}([(x, y, z)]) = T(x, y, z) = (x, y)$.

Back to de la Fuente:

Every real vector space X with dimension n is isomorphic to \mathbf{R}^n . What's the isomorphism?

Definition 4 Fix any Hamel basis $V = \{v_1, \dots, v_n\}$ of X . Any $x \in X$ has a unique representation

$$x = \sum_{j=1}^n \beta_j v_j$$

(here, we allow $\beta_j = 0$). Generally, vectors are represented as column vectors, not row vectors.

$$crd_V(x) = \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_n \end{pmatrix} \in \mathbf{R}^n$$

$crd_V(x)$ is the vector of coordinates of x with respect to the basis V .

$$crd_V(v_1) = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \quad crd_V(v_2) = \begin{pmatrix} 0 \\ 1 \\ \vdots \\ 0 \\ 0 \end{pmatrix} \quad crd_V(v_n) = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ 1 \end{pmatrix}$$

crd_V is an isomorphism from X to \mathbf{R}^n .

Matrix Representation of a Linear Transformation

Definition 5 Suppose $T \in L(X, Y)$, $\dim X = n$, $\dim Y = m$. Fix bases

$$\begin{aligned} V &= \{v_1, \dots, v_n\} \text{ of } X \\ W &= \{w_1, \dots, w_m\} \text{ of } Y \end{aligned}$$

$T(v_j) \in Y$, so

$$T(v_j) = \sum_{i=1}^m \alpha_{ij} w_i$$

Define

$$Mtx_{W,V}(T) = \begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & \vdots & \vdots \\ \alpha_{m1} & \cdots & \alpha_{mn} \end{pmatrix}$$

Notice that the columns are the coordinates (expressed with respect to W) of $T(v_1), \dots, T(v_n)$.

Observe

$$\begin{pmatrix} \alpha_{11} & \cdots & \alpha_{1n} \\ \vdots & & \vdots \\ \alpha_{m1} & \cdots & \alpha_{mn} \end{pmatrix} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} = \begin{pmatrix} \alpha_{11} \\ \vdots \\ \alpha_{m1} \end{pmatrix}$$

so

$$\begin{aligned} Mtx_{W,V}(T) \cdot crd_V(v_j) &= crd_W(T(v_j)) \\ \forall_{x \in X} Mtx_{W,V}(T) \cdot crd_V(x) &= crd_W(T(x)) \end{aligned}$$

Multiplying a vector by a matrix does two things:

- Computes the action of T
- Accounts for the change in basis

Example: $X = Y = \mathbf{R}^2$, $V = \{(1, 0), (0, 1)\}$, $W = \{(1, 1), (-1, 1)\}$,
 $T = id$, $T(x) = x$,

$$Mtx_{W,V}(T) \neq \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$Mtx_{W,V}(T)$ is the matrix which *changes basis* from V to W . How do we compute it?

$$v_1 = (1, 0) = \alpha_{11}(1, 1) + \alpha_{21}(-1, 1)$$

$$\begin{aligned}
\alpha_{11} - \alpha_{21} &= 1 \\
\alpha_{11} + \alpha_{21} &= 0 \\
2\alpha_{11} &= 1, \alpha_{11} = \frac{1}{2} \\
\alpha_{21} &= -\frac{1}{2} \\
v_2 = (0, 1) &= \alpha_{12}(1, 1) + \alpha_{22}(-1, 1) \\
\alpha_{12} - \alpha_{22} &= 0 \\
\alpha_{12} + \alpha_{22} &= 1 \\
2\alpha_{12} &= 1, \alpha_{12} = \frac{1}{2} \\
\alpha_{22} &= \frac{1}{2}
\end{aligned}$$

$$Mtx_{W,V}(id) = \begin{pmatrix} 1/2 & 1/2 \\ -1/2 & 1/2 \end{pmatrix}$$

Theorem 6 (3.5') *Let X and Y be vector spaces over the same field F , $\dim X = n$, $\dim Y = m$, bases*

$$\begin{aligned}
V &= \{v_1, \dots, v_n\} \text{ for } X \\
W &= \{w_1, \dots, w_m\} \text{ for } Y
\end{aligned}$$

Then

$$Mtx_{W,V} \in L(L(X, Y), F_{m \times n})$$

$Mtx_{W,V}$ is an isomorphism from $L(X, Y)$ to $F_{m \times n}$, the vector space of $m \times n$ matrices over F .

Theorem 7 ((From Handout)) *Let X, Y, Z be finite-dimensional vector spaces with bases U, V, W respectively, $S \in L(X, Y)$, $T \in L(Y, Z)$. Then*

$$Mtx_{W,V}(T) \cdot Mtx_{V,U}(S) = Mtx_{W,U}(T \circ S)$$

i.e. matrix multiplication corresponds via the isomorphism to composition of linear transformations. Note that $Mtx_{W,V}$ is a function from $L(X, Y)$ to the space of $m \times n$ matrices, while $Mtx_{W,V}(T)$ is an $m \times n$ matrix.

Proof: See handout. ■

The theorem can be summarized by the following “Commutative Diagram:”

$$\begin{array}{ccccc}
 & & S & & T \\
 & X & \rightarrow & Y & \rightarrow & Z \\
 crd_U & \updownarrow & & \updownarrow crd_V & & \updownarrow crd_W \\
 & \mathbf{R}^n & \rightarrow & \mathbf{R}^m & \rightarrow & \mathbf{R}^r \\
 & & Mtx_{V,U}(S) & & Mtx_{W,V}(T) & &
 \end{array}$$

We say the diagram commutes because you get the same answer any way you go around the diagram (in directions allowed by the arrows). The crd arrows go in both directions because crd is an isomorphism.

Section 3.5, Change of Basis, Similarity

Let X be a finite-dimensional vector space with basis V . If $T \in L(X, X)$ it is customary to use the same basis in the domain and range:

Definition 8

$$Mtx_V(T) \text{ denotes } Mtx_{V,V}(T)$$

Question: If W is another basis for X , how are $Mtx_V(T)$ and $Mtx_W(T)$ related?

$$Mtx_{V,W}(id) \cdot Mtx_W(T) \cdot Mtx_{W,V}(id)$$

$$\begin{aligned}
&= \text{Mtx}_{V,W}(id) \cdot \text{Mtx}_{W,V}(T \circ id) \\
&= \text{Mtx}_{V,V}(id \circ T \circ id) \\
&= \text{Mtx}_V(T) \\
&\text{Mtx}_{V,W}(id) \cdot \text{Mtx}_{W,V}(id) \\
&= \text{Mtx}_{V,V}(id) \\
&= \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & 1 & 0 & \cdots & 0 & 0 \\ & & & & & \\ & & & & & \\ 0 & 0 & 0 & \cdots & 0 & 1 \end{pmatrix} \\
\text{Mtx}_V(T) &= P^{-1} \text{Mtx}_W(T) P
\end{aligned}$$

where

$$P = \text{Mtx}_{W,V}(id)$$

is a change of basis matrix. On the other hand, if P is invertible, then P is a change of basis matrix (see handout).

Definition 9 Square matrices A, B are *similar* if

$$A = P^{-1}BP$$

for some invertible matrix P .

Theorem 10 *Suppose that X is finite-dimensional.*

- *If $T \in L(X, X)$ and U, W are any two bases of X , then $\text{Mtx}_W(T)$ and $\text{Mtx}_U(T)$ are similar.*
- *Conversely, given similar matrices A, B with $A = P^{-1}BP$ and any basis U , there is a basis W and $T \in L(X, X)$ such that*

$$\begin{aligned}
B &= \text{Mtx}_U(T) \\
A &= \text{Mtx}_W(T)
\end{aligned}$$

$$P = Mtx_{U,W}(id)$$

$$P^{-1} = Mtx_{W,U}(id)$$

Proof: See Handout on Diagonalization and Quadratic Forms. ■

Section 3.6: Eigenvalues and Eigenvectors

De la Fuente defines eigenvalues and eigenvectors of a matrix. Here, we define eigenvalues and eigenvectors of a linear transformation and show that λ is an eigenvalue of T if and only if λ is an eigenvalue for some matrix representation of T if and only if λ is an eigenvalue for every matrix representation of T .

Definition 11 Let X be a vector space and $T \in L(X, X)$. We say that λ is an *eigenvalue* of T and $v \neq 0$ is an *eigenvector corresponding to λ* if $T(v) = \lambda v$.

Theorem 12 (Theorem 4 in Handout) *Let X be a finite-dimensional vector space, and U any basis. Then λ is an eigenvalue of T if and only if λ is an eigenvalue of $Mtx_U(T)$. v is an eigenvector of T corresponding to λ if and only if $crd_U(v)$ is an eigenvector of $Mtx_U(T)$ corresponding to λ .*

Proof: By the Commutative Diagram Theorem,

$$T(v) = \lambda v \Leftrightarrow crd_U(T(v)) = crd_U(\lambda v)$$

$$\Leftrightarrow Mtx_U(T)(crd_U(v)) = \lambda(crd_U(v))$$

■ *Computing eigenvalues and eigenvectors:*

Suppose $\dim X = n$; let I be the $n \times n$ identity matrix. Given $T \in L(X, X)$, fix a basis U and let

$$A = Mtx_U(T)$$

Find the eigenvalues of T by computing the eigenvalues of A :

$$\begin{aligned}Av = \lambda v &\Leftrightarrow (A - \lambda I)v = 0 \\ &\Leftrightarrow (A - \lambda I) \text{ is not invertible} \\ &\Leftrightarrow \det(A - \lambda I) = 0\end{aligned}$$

We have the following facts:


- If $A \in \mathbf{R}_{n \times n}$,

$$f(\lambda) = \det(A - \lambda I)$$

is an n^{th} degree polynomial in λ with real coefficients; it is called the *characteristic polynomial* of A .

- f has n roots in \mathbf{C} , counting multiplicity:

$$f(\lambda) = (\lambda - c_1)(\lambda - c_2) \cdots (\lambda - c_n)$$

where $c_1, \dots, c_n \in \mathbf{C}$ are the eigenvalues; the c_j 's are not necessarily distinct. ** notice that $f(\lambda) = 0$ if and only if $\lambda \in \{c_1, \dots, c_n\}$, so the roots are the solutions of the equation $f(\lambda) = 0$.

- the roots which are not real come in conjugate pairs:

$$f(a + bi) = 0 \Leftrightarrow f(a - bi) = 0$$

- if $\lambda = c_j \in \mathbf{R}$, there is a corresponding eigenvector in \mathbf{R}^n .
- if $\lambda = c_j \notin \mathbf{R}$, the corresponding eigenvectors are in $\mathbf{C}^n \setminus \mathbf{R}^n$.

Diagonalization

Definition 13 Suppose X is finite-dimensional with basis U . Given a linear transformation $T \in L(X, X)$, let

$$A = Mtx_U(T)$$

We say that A can be diagonalized if there is a basis W for X such that $Mtx_W(T)$ is diagonal, i.e.

$$Mtx_W(T) = \begin{pmatrix} \lambda_1 & 0 & 0 & \cdots & 0 & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & \lambda_n \end{pmatrix}$$

Notice that the eigenvectors of $Mtx_W(T)$ are exactly the standard basis vectors of \mathbf{R}^n . But w_j is an eigenvector of T for λ_j if and only if $crd_W(w_j)$ is an eigenvector of $Mtx_W(T)$, and $crd_W(w_j)$ is the j^{th} standard basis vector of \mathbf{R}^n , so $W = \{w_1, \dots, w_n\}$ where w_j is an eigenvector corresponding to λ_j .

Then the action of T is clear: it stretches each basis element w_i by the factor λ_i .

Theorem 14 (6.7') *Let X be an n -dimensional vector space, $T \in L(X, X)$, U any basis of X , and $A = Mtx_U(T)$. Then the following are equivalent:*

- A can be diagonalized
- there is a basis W for X consisting of eigenvectors of T
- there is a basis V for \mathbf{R}^n consisting of eigenvectors of A

Proof: Use de la Fuente's Theorem 6.7 and Theorem 4 from the Handout. ■

Theorem 15 (6.8') *Let X be a vector space, $T \in L(X, X)$.*

- *If $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of T with corresponding eigenvectors v_1, \dots, v_m , then $\{v_1, \dots, v_m\}$ is linearly independent.*

- *If $\dim X = n$ and T has n distinct eigenvalues, then X has a basis consisting of eigenvectors of T ; consequently, if U is any basis of X , then $Mtx_U(T)$ is diagonalizable.*

Proof: This is an adaptation of the proof of Theorem 6.8 in de la Fuente. ■